

Proof Nets

A graph syntax for proofs

Reference:
Notes on proof-nets by Olivier Laurent

(Note: most slides are taken from the notes of Olivier Laurent)

1

MLL

$$\frac{}{\vdash A^\perp, A} ax \quad \frac{\vdash \Gamma, A \quad \vdash \Delta, A^\perp}{\vdash \Gamma, \Delta} cut$$

$$\frac{\vdash \Gamma, A \quad \vdash \Delta, B}{\vdash \Gamma, \Delta, A \otimes B} \otimes \quad \frac{\vdash \Gamma, A, B}{\vdash \Gamma, A \wp B} \wp$$

Recall that linear negation is defined :

$$(X^\perp)^\perp = X$$

$$(A \otimes B)^\perp = A^\perp \wp B^\perp$$

$$(A \wp B)^\perp = A^\perp \otimes B^\perp$$

2

Forgetting Sequential Structure

$$\frac{\frac{\frac{}{\vdash A, A^\perp} (ax) \quad \frac{}{\vdash B, B^\perp} (ax)}{\vdash A \otimes B, A^\perp, B^\perp} (\otimes) \quad \frac{}{\vdash C, C^\perp} (ax)}{\vdash (A \otimes B) \otimes C, A^\perp, B^\perp, C^\perp} (\otimes) \quad \frac{}{\vdash A, A^\perp} (ax) \quad \frac{}{\vdash B, B^\perp} (ax)}{\vdash (A \otimes B) \otimes C, A^\perp \wp B^\perp, C^\perp} (\wp) \quad \frac{}{\vdash A \otimes B, A^\perp, B^\perp} (cut)}{\vdash (A \otimes B) \otimes C, A^\perp, B^\perp, C^\perp} (\wp) \quad \frac{}{\vdash (A \otimes B) \otimes C, A^\perp \wp B^\perp, C^\perp} (\wp)}$$

3

Forgetting Sequential Structure

$$\frac{\frac{\frac{}{\vdash A, A^\perp} (ax) \quad \frac{}{\vdash B, B^\perp} (ax)}{\vdash A \otimes B, A^\perp, B^\perp} (\otimes) \quad \frac{}{\vdash C, C^\perp} (ax)}{\vdash (A \otimes B) \otimes C, A^\perp, B^\perp, C^\perp} (\otimes) \quad \frac{}{\vdash A, A^\perp} (ax) \quad \frac{}{\vdash B, B^\perp} (ax)}{\vdash (A \otimes B) \otimes C, A^\perp \wp B^\perp, C^\perp} (\wp) \quad \frac{}{\vdash A \otimes B, A^\perp, B^\perp} (cut)}{\vdash (A \otimes B) \otimes C, A^\perp, B^\perp, C^\perp} (\wp) \quad \frac{}{\vdash (A \otimes B) \otimes C, A^\perp \wp B^\perp, C^\perp} (\wp)}$$

4

Forgetting Sequential Structure

- (1) $\frac{}{\vdash A, A^\perp} ax$
- (2) $\frac{}{\vdash B, B^\perp} ax$
- (3) $\frac{}{\vdash A \otimes B, A^\perp, B^\perp} \otimes$
- (4) $\frac{}{\vdash C, C^\perp} ax$
- (5) $\frac{}{\vdash (A \otimes B) \otimes C, A^\perp, B^\perp, C^\perp} \otimes$
- (6) $\frac{}{\vdash (A \otimes B) \otimes C, A^\perp \wp B^\perp, C^\perp} \wp$
- (7) $\frac{}{\vdash A, A^\perp} ax$
- (8) $\frac{}{\vdash (A \otimes B) \otimes C, (A^\perp \wp B^\perp) \otimes A, C^\perp, A^\perp} \otimes$

5

Forgetting Sequential Structure

- (1) $\frac{}{\vdash A, A^\perp} ax$
- (2) $\frac{}{\vdash B, B^\perp} ax$
- (3) $\frac{}{\vdash A \otimes B, A^\perp, B^\perp} \otimes$
- (4) $\frac{}{\vdash C, C^\perp} ax$
- (5) $\frac{}{\vdash (A \otimes B) \otimes C, A^\perp, B^\perp, C^\perp} \otimes$
- (6) $\frac{}{\vdash (A \otimes B) \otimes C, A^\perp \wp B^\perp, C^\perp} \wp$
- (7) $\frac{}{\vdash A, A^\perp} ax$
- (8) $\frac{}{\vdash (A \otimes B) \otimes C, (A^\perp \wp B^\perp) \otimes A, C^\perp, A^\perp} \otimes$

6

Proof Nets

A graph syntax for proofs

7

Proof structures

A **proof structure** \mathcal{M} is a **labelled directed acyclic graph (DAG)** with possibly pending edges (i.e. some edges may have no source and/or no target) built over the alphabet of nodes which is represented below.
 (Note: in figures, the edges orientation is always top-bottom.)

- The **nodes** are labelled by ax, cut, \otimes , \wp
- The **edges** are labelled by MLL formulas.

For each node/link: **premises** = entering edges, **conclusions** = exiting edges

The **conclusions** of \mathcal{M} is the set of pending edges of \mathcal{M} .

8

In the graphical representation of a proof structure, we do not mention explicitly the direction of edges, but we draw them in such a way that direction is represented in a top-down way:

9

From proofs to proof structures

10

example

Translate each of these sequent calculus proofs. Start from axioms...

11

Is every structure the image of an MLL proof?

12

Proof Nets

A **PROOF NET** is a proof structure which is the image of an MLL proof

Internal condition!
Purely geometrical conditions (correction) characterize the proof structures which are proof nets

Theorem 1. A proof structure is correct iff it is a proof net.

Correctness criteria:

- (LT) Long trip [Girard]
- (AC) Connected-Acyclic [Danos-Regnier]
- ...

13

Danos-Regnier Criterion

Correctness Criterion

Switching Graphs

Correctness

- Switching graphs are **acyclic**.
- Switching graphs are **connected**.

14

15

Definition 2 (Correctness criterion AC (Danos-Regnier)). Let R be a proof structure.

A switching s is a function on the nodes of R , which chooses, for each \mathfrak{N} -link, either the left or the right premise.

A proof structure R is correct if for each switching, the unoriented graph obtained by erasing for each \mathfrak{N} -link of R the edges not chosen by s , is:

connected and acyclic

16

Acyclicity. A multiplicative proof structure is *acyclic* if its switching graphs do not contain any undirected cycle.

A proof structure with p \mathfrak{N} nodes induces 2^p switchings and thus 2^p switching graphs. A switching graph is not a proof structure in general since its \mathfrak{N} nodes have only one premise.

A *connected component* of a switching graph is a connected component of its underlying (undirected) multigraph.

17

Is this correct?

PN1:

PN2:

18

- Correctness guarantees:
 - ✓ Graph is image of a proof (sequentialization)
 - ✓ Normalization progresses (no deadlocks)
 - ✓ Normalization terminates (no infinite cycles)

19

Soundness

Proposition 4.1.1 (Soundness of Correctness). *The translation of a sequent calculus proof of MLL is a connected multiplicative proof net.*

20

Sequentialization

Theorem 4.1.1 (Sequentialization). *Any connected multiplicative proof net is the translation of a sequent calculus proof of MLL.*

21

Sequentialization answers the question:

We have a proof net. The problem: it is the image of a sequent calculus proof? And which?

22

The beauty of proof nets is normalization

23

Normalization
(local graph reductions!)

24

Let us try out!

$$\frac{\vdash A \wp A^\perp, A \otimes A^\perp \quad \vdash A^\perp \wp A}{\vdash A \wp A^\perp} \text{cut}$$

Write a proof net with this conclusion... and normalize it

25

How we write a proof net of these conclusions?

$A \wp A^\perp$ must type an edge conclusion of a par link, with premisses

$A \otimes A^\perp$ must type an edge conclusion of a tensor link, with premisses

Then we have to **choose** the axiom links!

26

Properties of normalization

Lemma (preservation of correctness)

If the proof structure R is correct and reduces to R' , then R' is correct.

27

Properties of normalization

1. Confluence?
2. Is normalization weakly/strongly normalizing?
3. Would you be able to define a normalizing strategy?
4. Would you be able to define a normalizing strategy which reaches normal form in a minimal number of steps?

28

Normalization of MLL proof-nets:

- Strongly normalizing
- Confluent
- Cut elimination: a proof-net in normal form contains no cuts

29

Let us try out!

$$\frac{\vdash A \wp A^\perp, A \otimes A^\perp \quad \vdash A^\perp \wp A}{\vdash A \wp A^\perp} \text{cut}$$

Write a proof net with this conclusion... and normalize it

30

• Let us try out another example

31

Let us try one more. First, write a proof net with this conclusion...

$$(X \otimes X) \multimap (X \otimes X) =$$

$$(X \otimes X)^\perp \wp (X \otimes X) = (X^\perp \wp X^\perp) \wp (X \otimes X)$$

TIP: How we write a proof net? As before, all proof nets with the same conclusion, start with the same nodes (the formula tree!)
What distinguishes different proofs are the axiom links

To distinguish the different occurrences of atoms, let us write indices:

$$(X_1^\perp \wp X_2^\perp) \wp (X_3 \otimes X_4)$$

In this case, we have two possible proofs, corresponding to two possible way to write axioms:
1,3 and 2,4
OR
1,4 and 2,3

32

In sequent calculus, they correspond to these two proofs (one uses exchange, one no)

$$\frac{\frac{\frac{\overline{\vdash X_3^\perp, X_1} \quad \overline{\vdash X_4^\perp, X_2}}{\vdash X_1^\perp, X_2^\perp, X_3 \otimes X_4} \wp}{\vdash X_1^\perp \wp X_2^\perp, X_3 \otimes X_4} \wp}{\vdash (X_1 \otimes X_3)^\perp \wp (X_2 \otimes X_4)} \wp$$

$$\frac{\frac{\frac{\overline{\vdash X_3^\perp, X_2} \quad \overline{\vdash X_4^\perp, X_1}}{\vdash X_1^\perp, X_2^\perp, X_3 \otimes X_4} \wp}{\vdash X_1^\perp \wp X_2^\perp, X_3 \otimes X_4} \wp}{\vdash (X_1 \otimes X_3)^\perp \wp (X_2 \otimes X_4)} \wp$$

33

When we have a formula whose normal proofs are exactly two, we have a good candidate to code BOOLEANS :

Let us indicate the formula $(X^\perp \wp X^\perp) \wp (X \otimes X)$ with B (for boolean).
We call one proof **true**, and the other **false**...

We can feed one of our two values to a proof which takes a boolean, and return a boolean.

We know that the normal form (i.e the result of computation) will be of type B... Hence one of our two values.

34

Try to normalize one of the proofs of $(X_1^\perp \wp X_2^\perp) \wp (X_3 \otimes X_4)$

with the proof net which has conclusions

$$(X_1 \otimes X_2) \otimes (X_3^\perp \wp X_4^\perp) \quad (X_5^\perp \wp X_6^\perp) \wp (X_7 \otimes X_8)$$


and axiom links: (1,6) (2,5) (3,7) (4,8)

What is the function coded by this proof net?

35

Correctness criterion, simplified

36



37

Correctness: if we focus on acyclicity, Danos-Regnier criterion can be reformulated (in equivalent way)

Let R be a proof structure; a *switching path* of R is a path which does not use any two edges entering on the same \exists link (such edges are called *switching edges*); a *switching cycle* is a switching path which is a cycle.

Definition 3 (Correctness criterion). A proof structure is correct if it does not contain any switching cycle.

$$\frac{}{\vdash A, A^\perp} (Ax) \quad \frac{\vdash \Gamma, A \quad \vdash \Delta, A^\perp}{\vdash \Gamma, \Delta} (Cut)$$

$$\frac{\vdash \Gamma, A \quad \vdash \Delta, B}{\vdash \Gamma, A \otimes B, \Delta} (\otimes) \quad \frac{\vdash \Gamma, A, B}{\vdash \Gamma, A \wp B} (\wp)$$


$$\frac{\vdash \Gamma \quad \vdash \Delta}{\vdash \Gamma, \Delta} (Mix)$$

we can throw away MIX later By requiring **connectness**


38

Exponentials


39



40



41



42

What is the associated proof-net?

$$\frac{\frac{\frac{}{\vdash A, A^\perp} ax}}{\vdash A \wp A^\perp} \wp}{\vdash A \wp A^\perp, ?B} ?w \quad \mapsto$$

$$!A \otimes !A \dashv\vdash !A$$

$$?A \wp ?A \dashv\vdash ?A$$


Can you write the proof so that all axioms are atomic (ie on atomic formulas)?

43

Reduction steps


44

Reduction Steps: ?d




45

Reduction Steps: ?w




46

Reduction Steps: ?c




47

- étape 1/?c :



48

Reduction Steps: ?p



49

Confluence?
Weak Normalization?
Strong Normalization?

50

Properties of MELL reduction

1. Is confluent?
2. Is weakly normalizing?

Tip for WN.

Given a proof-net R, try to make decrease a size S(R).

For example:

- **Size of a cut:** pair (s,t) where s is the size of the cut and, t is the size of the ?-tree above the ? premisses of the cut if any, or 0
- **Size S(R) of the proof-net R:** the multiset of the sizes of all its cuts


51

Weak Normalization

52

Weak Normalization

- cut relation: $c_1 \prec c_2$ (exponential cuts only)



- correctness \implies no cycle in \prec^* \implies maximal cuts
- reduction of a maximal cut \implies [([formula], |?-tree|)] decreases

53

Weak Normalization

Size of a cut: pair (s,t) where s is the size of the cut and, t is the size of the ?-tree above the ? premisses of the cut if any, or 0


Size S(R) of the proof-net R: the multiset of the sizes of all its cuts

The size of a ?-tree is its number of nodes.
descent path (bis): from a node downwards to a conclusion or to a cut or to a premisses of ! node (that is we do not continue down through an ! node)


54

Confluence


55




56




57



58



59



60

Bonus Exercise

A proof-net is *polarized* if every edge is labelled by a positive or a negative formula

Let M be a $MELL$ MLL polarized proof structure. We denote by $Pol(M)$ the graph which has the same nodes and edges as M , but where the edges are directed downward if positive, upwards if negative.

Do you see any *simple* way to show that the following are equivalent?


- (1.) M is acyclic correct, (2.) $Pol(M)$ is a DAG.

61

Sequentialization

62


We have a proof net. The problem: it is the image of a sequent calculus proof? And which?



63


In fact, what is a sequent calculus proof? A sequent calculus proof is a tree of rules...

$$\frac{\frac{\frac{}{\vdash A, A^\perp} ax \quad \frac{}{\vdash B, B^\perp} ax}{\vdash A^\perp \otimes B^\perp, A, B} \otimes}{\vdash A^\perp \otimes B^\perp, A \wp B} \wp \quad \frac{}{\vdash C, C^\perp} ax}{\vdash A^\perp \otimes B^\perp, (A \wp B) \otimes C^\perp, C} \otimes \quad \frac{}{\vdash D, D^\perp} ax}{\vdash A^\perp \otimes B^\perp, (A \wp B) \otimes C^\perp, C \otimes D, D^\perp} \otimes$$



64

Such a tree directly corresponds to the following sequent calculus proof:



$$\frac{\frac{\frac{}{\vdash A, A^\perp} ax \quad \frac{}{\vdash B, B^\perp} ax}{\vdash A^\perp \otimes B^\perp, A, B} \otimes}{\vdash A^\perp \otimes B^\perp, A \wp B} \wp \quad \frac{}{\vdash C, C^\perp} ax}{\vdash A^\perp \otimes B^\perp, (A \wp B) \otimes C^\perp, C} \otimes \quad \frac{}{\vdash D, D^\perp} ax}{\vdash A^\perp \otimes B^\perp, (A \wp B) \otimes C^\perp, C \otimes D, D^\perp} \otimes$$

65

Translating lambda-calculus into LL

CbN and CbV

66

Simply typed lambda calculus

$$\frac{}{\Gamma, x : \tau \vdash x : \tau} \text{ (var)} \quad \frac{\Gamma, x : \tau \vdash t : \sigma}{\Gamma \vdash \lambda x.t : \tau \rightarrow \sigma} \text{ (abs)} \quad \frac{\Gamma \vdash t : \tau \rightarrow \sigma \quad \Gamma \vdash u : \tau}{\Gamma \vdash tu : \sigma} \text{ (app)}$$

$$X^* = X \qquad X^* = !X$$

$$(A \rightarrow B)^* = !A^* \multimap B^* \qquad (A \rightarrow B)^* = !(A^* \multimap B^*)$$

$$(\Gamma \vdash A)^* = !\Gamma^* \vdash A^* \qquad (\Gamma \vdash A)^* = \Gamma^* \vdash A^*$$

67

CbN translation

$$\frac{}{\Delta, x : A \vdash x : A} \text{ (var)} \quad \frac{\Gamma \vdash t : \tau \rightarrow \sigma \quad \Gamma \vdash u : \tau}{\Gamma \vdash tu : \sigma} \text{ (app)}$$

$$\frac{\Gamma, x : \tau \vdash t : \sigma}{\Gamma \vdash \lambda x.t : \tau \rightarrow \sigma} \text{ (abs)}$$

68

Three lambda calculi... Or indeed just one.

The **two Girard's translations** (intuitionistic into linear logic) are well-known to correspond to the **CbN** and **CbV**.

Taking this point of view provides a modular approach.

In $\Lambda_{\oplus}^!$ the natural constraint is:

- no reduction in the scope of a !-box

Girard's translations transform this policy respectively in:

- $\Lambda_{\oplus}^{\text{cbv}}$: no reduction under λ -abstraction
- $\Lambda_{\oplus}^{\text{cbn}}$: no reduction argument position.

69

linear λ -calculus

$$\Lambda_{\oplus}^!$$

$\Lambda_{\oplus}^{\text{cbv}}$ $\xrightarrow{\text{Girard Translation Cbv}}$ $\Lambda_{\oplus}^!$ $\xleftarrow{\text{Girard Translation Cbn}}$ $\Lambda_{\oplus}^{\text{cbn}}$

CbV λ -calculus **CbN λ -calculus**

The translation gives :

- Confluence
- Standardization (finite and asymptotical)

70

CbV translation

$$X^* = !X$$

$$(A \rightarrow B)^* = !(A^* \multimap B^*)$$

$$(\Gamma \vdash A)^* = \Gamma^* \vdash A^*$$

71

CbV

72